

The University of Chicago

THE BIURET REACTION OF THE PENTAPEPTIDE TETRAGLYCYLGLYCINE

A PART OF A DISSERTATION SUBMITTED TO THE
FACULTY OF THE DIVISION OF THE PHYSICAL
SCIENCES IN CANDIDACY FOR THE DEGREE OF
DOCTOR OF PHILOSOPHY

DEPARTMENT OF CHEMISTRY

1934

By
PAUL E. WENAAS

Private Edition, Distributed by
THE UNIVERSITY OF CHICAGO LIBRARIES
CHICAGO, ILLINOIS

[Reprinted from the Journal of the American Chemical Society, **59**, 1353 (1937).]

The University of Chicago

THE BIURET REACTION OF THE PENTAPEPTIDE TETRAGLYCYLGLYCINE

A PART OF A DISSERTATION SUBMITTED TO THE
FACULTY OF THE DIVISION OF THE PHYSICAL
SCIENCES IN CANDIDACY FOR THE DEGREE OF
DOCTOR OF PHILOSOPHY

DEPARTMENT OF CHEMISTRY

1934

By
PAUL E. WENAAS

Private Edition, Distributed by
THE UNIVERSITY OF CHICAGO LIBRARIES
CHICAGO, ILLINOIS

[Reprinted from the Journal of the American Chemical Society, **59**, 1353 (1937).]




The Biuret Reaction of the Pentapeptide Tetraglycylglycine¹

BY PAUL E. WENAAS

As a continuation of the study of the biuret reaction of the polypeptides of the glycine series in preparation for a study of the biuret reaction of the proteins,² the biuret salt of the pentapeptide tetraglycylglycine, $\text{NH}_3^+(\text{CH}_2\text{CONH})_4\text{CH}_2\text{CO}_2^-$, has been isolated and examined.

The biuret salt of tetraglycylglycine was prepared by the saturation with cupric hydroxide of a solution of the pentapeptide in aqueous sodium hydroxide. The analytical data for the completely dehydrated salt obtained from this aqueous solution by precipitation with an absolute alcohol-absolute ether mixture agree quite favorably with the empirical formula $[\text{Na}_3^+\text{Cu}^{++}(\text{NH}_2(\text{CH}_2\text{CO}^-\text{=N})_4\text{CH}_2\text{CO}^-)]$ for the salt. The amount of sodium found, however, was lower than that calculated by a greater amount than would result from any possible analytical errors, and at the same time the hydrogen found was correspondingly higher than that required by theory. These results suggest two possibilities: (1) that the trisodium salt was hydrolyzed in part during its preparation, or (2) that not all the acid hydrogen atoms in the pentapeptide molecule were neutralized by the alkali used. The molecular weight of the salt was not determined. If its molecular and empirical formulas are identical the structure of the salt may be



The structure assigned may be interpreted as follows: the enol tautomer of tetraglycylglycine is considered to contain five ionizable hydrogen atoms. In the course of the formation of a molecule of sodium copper tetraglycylglycine three

of these are considered to be neutralized by the alkali used. When the acid strength of the peptide molecule is thus reduced by the neutralization of three acid groups, the amine character of the peptide is enhanced and cupric ion then may enter a complex of the cupri-ammonium type with four of the strongly basic nitrogen atoms of the peptide molecule. The two remaining acid groups of the enolate serve as the negative radical of the "complex salt," as sulfate ion is the negative ion of copper ammonium sulfate. The term "complex salt" as here used signifies that part of the biuret salt molecule which is placed within the brackets in the structure given. The negative radical of the cupri-peptide complex is not free to ionize, since this radical is an integral part of the peptide molecule.

Experimental

Synthesis of Peptides.—The synthesis of tetraglycylglycine was accomplished according to the procedure described by Fischer.³ Glycine was converted into its ester hydrochloride from which was obtained glycine anhydride. The anhydride yielded upon hydrolysis and treatment with chloroacetyl chloride in one operation chloroacetylglycylglycine, which was converted by the use of aqueous ammonia into diglycylglycine. The treatment of the tripeptide successively with chloroacetyl chloride and ammonia yielded triglycylglycine, which on treatment successively with chloroacetyl chloride and aqueous ammonia gave tetraglycylglycine.

Sodium Copper Tetraglycylglycine, $\text{Na}_3\text{CuC}_{10}\text{H}_{12}\text{N}_6\text{O}_6$.—The salt was prepared as follows: a solution of 0.1 g. of tetraglycylglycine in 7.16 ml. of water was treated with 2.84 ml. of a 1.754% solution of carbonate free sodium hydroxide. The resultant solution was shaken with an excess of cupric hydroxide for fifteen minutes.⁴ After removal of the excess cupric hydroxide on a Neubauer micro-platinum filtering crucible having a porous iridium bottom, the biuret salt was precipitated from the reaction mixture by the addition of 140 ml. of absolute ethyl alcohol and 100 ml. of absolute ether. The salt was collected, dried at room temperature over phosphorus pentoxide and analyzed.

Anal. Calcd. for $\text{Na}_3\text{CuC}_{10}\text{H}_{12}\text{N}_6\text{O}_6$: Na, 16.02; Cu,

(1) The work here described, conducted under the direction of Dr. Mary Rising Stieglitz, constitutes part of the dissertation of the writer, presented in partial fulfillment of the requirements for the doctorate degree at the University of Chicago.

(2) Rising, Parker and Gaston, *J. Am. Chem. Soc.*, **56**, 1178 (1934).

(3) E. Fischer, "Untersuchungen über Aminosäuren, Polypeptide und Proteine," Vols. I and II, Verlag von Julius Springer, Berlin, 1906.

(4) The cupric hydroxide was prepared by the method described by Böttger, *J. prakt. Chem.*, **73**, 491 (1858).

14.76; N, 16.26; C, 27.86; H, 2.81. Found: Na, 13.20, 13.62; Cu, 14.35, 14.85; N, 15.47, 15.55; C, 27.14, 27.30; H, 3.73, 3.69.

The salt was dehydrated completely at 78° over phosphorus pentoxide. A sample of the salt weighing 25.270 mg. dried at room temperature over phosphorus pentoxide when completely dehydrated at 78° lost 0.378 mg. weight, or 1.50%. This dehydrated salt was analyzed for nitrogen.

Anal. Calcd. for $\text{Na}_3\text{CuC}_{10}\text{H}_{12}\text{N}_5\text{O}_6$: N, 16.26. Found: N, 15.81, 15.98.

The analytical data of the product dried at room temperature were corrected for the loss of weight on complete dehydration at 78°.

Anal. Calcd. for $\text{Na}_3\text{CuC}_{10}\text{H}_{12}\text{N}_5\text{O}_6$: Na, 16.02; Cu,

14.76; N, 16.26; C, 27.86; H, 2.81. Found (corr.): Na, 13.41, 13.83; Cu, 14.57, 15.07; N, 15.71, 15.79; C, 27.56, 27.72; H, 3.62, 3.58.

Sodium copper tetraglycylglycine is a hygroscopic pink powder. When heated in a capillary-size melting point tube it begins to decompose with a blackening of the sample at 268°, shrinks to a considerable extent at 272°, and evolves gas at 279–281° (all temp. corr.).

Summary

The sodium biuret salt of the pentapeptide tetraglycylglycine has been isolated and analyzed. The character of this salt is discussed in terms of the theory of the biuret reaction so far developed.

